CREATIVE BIOMART®
Assay Kit

TR-FRET CAR Coactivator Assay Kit, rabbit

Product Information

Cat

Kit-0892

Common Name

TR-FRET

Cat.No. Kit-0892

Description

The TR-FRET Constitutive Androstane Receptor (CAR) Coactivator Assay provides a sensitive and robust method for highthroughput screening (HTS) of potential CAR ligands as agonists of ligand-dependent coactivator recruitment or inverse agonists of liganddependent coactivator displacement. The kit uses a terbium-labeled anti-GST antibody, a fluorescein-labeled coactivator peptide, and a CAR ligand-binding domain (CAR-LBD) that is tagged with glutathione-S-transferase (GST) in a homogeneous mix-and-read assay format.
To assay:
When running the TR-FRET CAR Coactivator Assay, CAR-LBD is added to ligand test compounds followed by addition of a mixture of the fluorescein-coactivator peptide and terbium anti-GST antibody. After an incubation period at room temperature, the 520 nm/ 495 nm TR-FRET ratio is calculated and can be used to determine the EC50 from a dose response curve of the compound. Based on the biology of the CAR-coactivator peptide interaction, this ligand's EC50 is a composite value representing the amount of ligand required to bind to receptor, effect a conformational change, and either recruit or displace coactivator peptide.

Applications

Nuclear Receptor Assay

Storage

Tb-anti-GST Antibody (Rabbit): Store at -20°C
CAR-LBD, GST: Store at -80°C
Fluoroscein Coactivator Peptide: Store at -20°C

Tel: 1-631-559-9269 1-516-512-3133
Email:info@creative-biomart.org
Fax:1-631-938-8127
45-1 Ramsey Road, Shirley, NY 11967, USA

CREATIVE BIOMART®
Assay Kit

TR-FRET CAR Coactivator Assay Kit, rabbit

TR-FRET Coregulatore Buffer G: Store at 4°C
1M DTT: Store at -20 °C or -80 °C

Size

800 x 20 µL assays

Materials Required but Not Supplied

Microplate Reader

Detection method Fluorescent

Tel: 1-631-559-9269 1-516-512-3133

Email:info@creative-biomart.org

Fax:1-631-938-8127

45-1 Ramsey Road, Shirley, NY 11967, USA