CREATIVE BIOMART®
Assay Kit

SIRT2 fluorometric drug discovery assay Kit

Product Information

Cat.No. Kit-0786

Product Overview

Useful for inhibitor screening or characterizing enzyme kineticsIncludes optimal substrate selected from a panel of acetylated sites in p53 and histonesSupplied with enough recombinant enzyme for 96 assays (1 x 96-well plate)

Size

96 wells

Description

A FLUOR DE LYS fluorescent assay system. The SIRT2 Fluorescent Activity Assay/Drug Discovery Kit is a complete assay system designed to measure the lysyl deacetylase activity of the recombinant human SIRT2 included in the kit. The kit is ideal for chemical library screening for candidate inhibitors or activators or kinetic assay of the enzyme under varying conditions. The FLUOR DE LYS SIRT2 assay is based on the FLUOR DE LYSSubstrate and FLUOR DE LYS Developer combination. The assay procedure has two steps. First, the FLUOR DE LYS SIRT2 Substrate, which comprises a unique peptide based on amino acids 317-320 of p53 (Gln-Pro-Lys-Lys (Ac)), is incubated with SIRT2. Deacetylation of the substrate sensitizes the substrate so that, in the second step, treatment with the FLUOR DE LYSDeveloper II produces a fluorophore.

Applications

Fluorescence microscopy, HTS

Storage

-80°C

Kit Components

SIRT2 (Sirtuin 2, hSir2) (human, recombinant)(500 U, one U=1 pmol/min at 37°C, 500μM, FLUOR DE LYS-SIRT2 substrate, 500μM NAD ; Recombinant enzyme dissolved in 25mM TRIS, pH 7.5, 100mM sodium chloride, 5mM dithiothreitol and 10% glycerol. See vial label for activity and protein

Tel: 1-631-559-9269 1-516-512-3133
Email:info@creative-biomart.org
Fax:1-631-938-8127
45-1 Ramsey Road, Shirley, NY 11967, USA

CREATIVE BIOMART®
Assay Kit

SIRT2 fluorometric drug discovery assay Kit

concentrationsStorage: -70°C, avoid freeze/thaw cycles!FLUOR DE LYS-SIRT2, Deacetylase substrate(100µl; 5mM solution in 25mM TRIS/Cl, pH 8.0, 137mM sodium chloride, 2.7mM potassion chloride, 1mM magnesium chloride)FLUOR DE LYS Developer II Concentrate (5x) (5 x 250 µl; 5x Stock Solution; Dilute in Assay Buffer before useStorage: -70°CNAD+ (Sirtuin Substrate)(500 µl; 50 mM β-Nicotinamide adenine dinucleotide (oxidized form) in 50mM TRIS/CL, pH 8.0, 137mM sodium chloride, 2.7mM potassium chloride, 1mM magnesium chloride)Storage: -70°CNicotinamide (Sirtuin Inhibitor)(500µl; 5 mM Nicotinamide in 50mM TRIS/Cl, pH 8.0, 137mM sodium chloride, 2.7mM potassium chloride, 1mM magnesium chloride)Storage: -70°CSuramin sodium (Sirtuin Inhibitor)(10 mg; Solid MW: 1429.2, soluble in water or assay buffer (to 25mM))Storage: -70°CFLUOR DE LYS Deacetylated Standard (30 µl; 10mM in DMSO)Storage: -70°CSirtuin Assay Buffer(20 ml; 50mM TRIS/Cl, pH 8.0, 137mM sodium chloride, 2.7mM potassium chloride, 1mM magnesium chloride, 1 mg/ml bovine serum albumin)Storage: -70°C1/2 volume microplateStorage: Room temperature1/2 volume white microplateStorage: Room temperature

Tel: 1-631-559-9269 1-516-512-3133
Email:info@creative-biomart.org
Fax:1-631-938-8127
45-1 Ramsey Road, Shirley, NY 11967, USA